

## COMPARATIVE STUDY OF LOCAL CONCENTRATE WITH IMPORT CONCENTRATE INCLUSION IN BROILER CHICKS DIETS

Mukhtar Ahmed Mukhtar and Mohamed H. Tabidi

*Department of Animal Production, College of Agricultural Studies, Sudan University of Science and Technology (sust.edu)*

**ABSTRACT:** A total of 120 broiler chicks, were used in this experiment to evaluate the effect of feeding local concentrate replaced the imported concentrate on the performance of broiler chicks. Three dietary treatments were formulated as: diet (A) containing 5% imported concentrate (IC), diets B and C containing 5% and 7.5% local concentrate (LC) respectively. The experiment was replicated four times with 10 chicks / replicate. The experiment was lasted for six weeks. The measured parameters were, average body weight, average feed intake average body weight gain and feed conversion ratio. Results obtained revealed that substitution of imported concentrate with local concentrate had significantly ( $P<0.05$ ) positive effect on chicks performance. Whereas, chicks fed diet with 7.5% LC significantly observed best performance. However, chicks fed on 5% LC showed significantly ( $P<0.05$ ) the best FCR. Substitution of local concentrate (LC) with LC economically profitable.

**KEYWORDS:** concentrate, Substitution, chick's performance.

### INTRODUCTION

In Sudan, concentrates have been used till now in poultry production due to its vital role to complete the protein and microelements in poultry feeds so, to maximize the growth performance of birds. Today the poultry industry, in Sudan faced, feed crisis because of high cost of production which attributed to raise of cost of feed ingredients mainly imported concentrates ([Mukhtar et al., 2010](#)). Now there were many attempts from nutritionists to replace the imported concentrate with different locally available protein sources. [Omer, \(2001\)](#) and [Mukhtar et al. \(2010\)](#) completely replaced imported concentrate by synthetic lysine and methionine and they recorded significantly improvement on the chick's performance. Fishmeal is a natural balanced feed ingredient that is high in protein, energy, minerals (Calcium and Phosphorus), natural source of vitamins (including choline, biotin and vitamin B<sub>12</sub>, A and E) and the microelements, selenium and iodine. The protein in fishmeal has a highly biological value in diets for animals ([Pike, 1999](#)). It is rich in the essential amino acids particularly lysine and the sulfur amino acids, ([Anonymous, 2002](#)). Fishmeal naturally contains from 62% to 64% proteins and digestible fats. Its quality varies due to the variety, parts of fish and processing technologies ([Fickler, 2002](#); [Dale et al., 2004](#)). The objective of this study is to evaluate the completely substitution of imported concentrate

with locally produced one on the performance of broiler chicks and cost of production.

### MATERIALS AND METHODS

One hundred and twenty, day-old, unsexed broiler chicks (Ross 308) were brooded at the teaching and Research Farm of College of Agricultural Studies, Sudan University of Science and Technology. Animals were fed with 5% and 7.5% local concentrate for the first week and control diet, respectively. The chicks were assigned to the three experimental diets in a complete randomized design, 40 chicks per treatment group, each treatment groups was further subdivided into four replicates of 10 chicks. Feed and water were supplied *ad-libitum*, and continuous light was provided throughout experimental period. Chicks were vaccinated against Newcastle and Gumboro diseases. Vitamins were offered before and after vaccinations. Measurements taken were, feed intake, live body weight gain, feed conversion ratio (FCR) and mortality rate. At the end of 6 weeks experiment, 4 chicks from each treatment were randomly selected after overnight fast (except for water), weighted individually, slaughtered, carcasses were scaled in hot water and de-feathered by hand, then eviscerated. The eviscerated carcasses were stored in refrigerator under -4°C for 24 hours, then they were divided into two halves, each half was dissected into commercial cuts (Thigh, breast, drumstick) carefully from their points of origin

and weighted individually and each one expressed as the percent of hot carcass.

Data collected were subjected to analysis of variance; means were separated using [Duncan's Multiple Range Test \(1955\)](#).

**Table 1:** Percent of ingredients inclusion of experimental diets


Ingredients	Control(5%IC)	5%LC	7.5%LC
Sorghum	65.75	64.142	62.1
Groundnut cake	13.0	14.0	13.0
Sesame cake	15.0	15.0	15.0
Imported concentrate	5.0	-	-
Local concentrate	-	5.0	7.5
Oyster	1.0	0.487	1.95
Salt	0.25	0.25	0.25
Lysine	-	0.344	-
Methionine	-	0.159	0.0

## RESULTS

The effect of feeding by different levels of local concentrate (LC) on broiler performance was summarized in Table (2) and (Figure 1). Chicks with diets containing 5% or 7.5% local concentrate had significantly ( $P<0.05$ ) higher body weight and average body weight gain compared with chicks fed diet containing 5% local concentrate consumed significantly ( $P<0.05$ ) low feed compared with those fed on 5% imported concentrate and 7.5% local concentrate at the same time recorded significantly ( $P<0.05$ ) the best feed conversion ratio. However, broiler final body weight average body weight, feed intake and feed conversion ratio were increasing improved by increasing the level of local concentrate (7.5%) inclusion to the diet. The health of the experimental chicks did not affect with the type of the concentrate, however, chicks fed on 7.5% local concentrate recorded zero mortality. Chicks fed with diet containing local concentrate (5% and 7.5%) were recorded more revenue and profit compared to chicks fed with imported concentrate although, chicks fed with 5% local concentrate observed the highest revenue and profit.

**Table 2:** Effect of feeding broiler chicks with different local concentrate compared with imported concentrate

Parameters	Control(5%IC)	5%LC	7.5%LC
Initial weight	213.5	228.7	218.5
Final weight	1930.8	2071	2078.5
Feed intake	3938.1	2971.84	3632.9
Body weight gain	1717.3	1842.3	1860
FCR	2.0	1.61	1.95



**Figure 1:** Effect of feeding broiler chicks with different local concentrate compared with imported concentrate

**Table 3:** Economical efficiency of feeding broiler chicks with different local concentrate compared with imported concentrate

Items	Control (5%IC)	5% LC	7.5% LC
Cost	13.36	14.29	14.15
Income	15.64	17.94	17.25
Profit	2.28	3.62	3.1
Profitable	1.0	1.59	1.36

## DISCUSSION

The local concentrate contained high level of fishmeal (20%) with synthetic amino acids mainly synthetic lysine (12%) and methionine (4.3%) and plant protein sources (groundnut cake and sesame cake).

Chicks fed on different local concentrate performed significantly better compared to chicks fed on diet containing local concentrate. They recorded 7.26% of 7.65% more body weight, and 7.28%, 8.3% for body weight gain. Chicks fed on 5% and 75% LC respectively.

This might be due to the high level of protein from many sources and with high biological value and to adequate higher available amino acids. These results were in line with the result of [Karimi, \(2006\)](#) who recorded that fishmeal improved broiler performance and the improvement affected with the level of fishmeal in the diet. Also these results were in agreement with the findings of [Fritts et al., \(2000\)](#); [Omer, \(2001\)](#) and [Mukhtar et al., \(2010\)](#) who found that the broiler chicks performance improved due to adequate of higher available of amino acids. On contrary [Agbede and Aleter, \(2003\)](#) who replaced fishmeal with leaf protein contrite from glycidin which caused significant reduction in the broiler chicks performance.

Mortality of experimental chicks was within normal range although chicks fed diets with 7.5% local concentrate recorded no mortality, this might be due to that fishmeal protect health and welfare, results were agreed with that of ([Anonymous, 2002](#)).

**REFERENCES**

- Agbede JO, Aletor VA. Evaluation of Fish Meal Replaced with Leaf Protein Concentrate from *Glyricidia* in Diets for Broiler - Chicks: Effect on Performance, Muscle Growth, Haematology and Serum Metabolites. *International Journal of Poultry Science* 2003; 2(4): 242-250.
- Anonymous. Fishmeal facts and figures. Fishmeal information Network (FIN). International fishmeal and oil manufactures association 2002.
- Dale NM, Zumbado M, Gernat AG, Romo G. Nutritional value of tilapia meal. *J Appl Poult Res* 2004;13:370-372.
- Duncan DB. Duncan's new multiple range test. *Metric Approach* 1955;11:1-42.
- Fickler J. Fish meal: high protein does not stand for high quality. *Feed Int* 2002;23:13-16.
- Fritts CA, Motl MA, Si J, Waldroup PW. Interaction of lysine and methionine in diets for growing broilers. *Poult Sci* 2000;79(1): 128.
- Karimi A. The effect of varying fishmeal inclusion levels (%) on performance of broiler chicks. *International Journal of Poultry Science* 2006; 5(3): 255-258.
- Mukhtar MA, Mohammed KA, Musa MH. Replacement Value of Lysine and Methionine for Super Concentrate in Broiler Chick's Yield and Quality. *Journal of Science and Technology* 2010;11(2): 28-29.
- Omer MA. Replacement of Super-Concentrate by L-lysine and DL-methionine at Different Levels in Broiler Rations. MSc. Thesis, Sudan University of Science and Technology, Sudan 2001.
- Pike IH. Health benefits from feeding fish oil and fish meal: The role of long chain omega-3 polyunsaturated fatty acids in animal feeding. *International fishmeal and oil manufactures association* 1999;28.